# Budgerigars

Instructor: Sohail Ahmad

# Budgerigars





#### **Budgerigars**

- Budgerigars are the smallest type of birds' resembling with parrots;
  also called as psittacine, parakeets, passerine and budgie
- These are the Australian parrots because their origin is Australia
- Mostly present in southern part of Australia
- These are very old birds; native to Australia

#### Why they called budgerigars?

- Budgerigars, in old local language of Australia budgeri means good and gars means cockatoo (parrots); so, budgerigars mean good looking parrots
- Different types and colors of budgerigars are found in zoo and aviaries


#### Classification of Budgerigars

Kingdom Animalia Phylum Chordata Sub-phylum Vertebrata Class Aves **Psittaciformes** Order Psittacidae Family Genus Melopsittacus Specie Undulates


The zoological name of budgerigars is *Melopsittacus undulates* 

- The scientific name of budgerigars is Melopsittacus undulates
- Melo means melodious bird; psittacus is the class of parrots and undulates means the article or thing having some movements
- Birds give movement to their feathers and tails

- Usually we keep these birds at our home
- Must be care when feeding the birds on the cages because if the birds having infection it may cause a respiratory disease in humans called as psittacosis







If maintaining parrots, it is required to take care of your birds, try to sprinkle water on cage so that fecal material may not be dry

#### Precautions while maintain birds

- Try to don't handle fecal material of bird directly
- When entering the cage don't take or smell birds
- Mostly wear gloves
- After feeding the birds wash your hands with good disinfectant or soap

### **Body weight**

- Its size is very small lesser than quail
- Mature body weight is just 30-40 grams
- Body weight of day old chick is less than 2 grams about 1.5-2.0 grams
- Total length of bird is 7 inches in some cases it may go up to 8 inches

#### Pre-cocial birds

- Bird species whose chick are hatched in an advanced state and are able to feed them almost immediately are called as pre-cocial birds
- Have fully developed feathers that help in maintaining body temperature
- They are not dependent on their parents

#### Altricial birds

- Birds whose day old chicks are naked called as altricial birds
- Their feathers are not developed at the time of hatching
- They can't maintain their body temperature
- Their mother makes fully arrangement during early life

#### **Incubation period**

Incubation period is almost 18 day; It may range from 18-20 days

#### **Fledgling time**

■ The time when bird is mature and able to take flight is called as fledgling time; Fledgling time in budgerigars is 4-5 weeks

#### Sex ratio

Sex ratio in budgerigars is 1:1; mostly in pair

#### Size of cage

For one pair of budgerigars we make 18×18 inches of cage whose length is 30 inches, because bird has ability to take flight

#### Natural incubation

- In natural incubation, female like to lay egg in the nest box
- Number of eggs is 4-8 sometimes it may go up to 10 eggs in a clutch
- There may be few clutches in a season

#### **Breeding season**

- Breeding season in budgerigars is rainy season when climate is moderate because birds are very sensitive
- They are herbivores or grainivores
- 5-6 million birds are present in the world

#### Sex distinguish

- Sex distinguish is from cero (nasal area) check
- In adult male color of cero is blue while in case of female it is brown or mixture of brown
- There are different colors of budgerigars from which most common colors are yellow, green, and white



#### **Color Pattern**

- **■**Green Series
- **Blue Series**

# Light Green



### **Dark Green**



### Olive Green



## Lutino



# Sky blue



### Mauve



## Albino



# Grey



## **Violet**

